

# Anti-C/EBP delta/epsilon Antibody

Catalog # AP53659

## Product Information

---

<b>Application</b>	WB, IF
<b>Primary Accession</b>	<a href="#">P49716</a>
<b>Other Accession</b>	<a href="#">Q15744</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	28467

## Additional Information

---

<b>Gene ID</b>	1052
<b>Other Names</b>	CEBPD; CCAAT/enhancer-binding protein delta; C/EBP delta; Nuclear factor NF-IL6-beta; NF-IL6-beta; CEBPE; CCAAT/enhancer-binding protein epsilon; C/EBP epsilon
<b>Target/Specificity</b>	Recognizes endogenous levels of C/EBP delta/epsilon protein.
<b>Dilution</b>	WB~~1/500 - 1/1000 IF~~1/50 - 1/200
<b>Format</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information


---

<b>Name</b>	CEBPD
<b>Function</b>	Transcription activator that recognizes two different DNA motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers (PubMed: <a href="#">16397300</a> ). Important transcription factor regulating the expression of genes involved in immune and inflammatory responses (PubMed: <a href="#">16397300</a> , PubMed: <a href="#">1741402</a> ). Transcriptional activator that enhances IL6 transcription alone and as heterodimer with CEBPB (PubMed: <a href="#">1741402</a> ).
<b>Cellular Location</b>	Nucleus.


## Background

---


Rabbit polyclonal antibody to C/EBP delta/epsilon



Western blot analysis of C/EBP delta/epsilon expression in CT26 (A), HeLa (B) whole cell lysates.



Immunohistochemical analysis of C/EBP delta/epsilon staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of C/EBP delta/epsilon staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.