

Anti-CHRNA7 Antibody

Catalog # AP53661

Product Information

Application	WB
Primary Accession	P36544
Other Accession	Q494W8
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Bovine, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56449

Additional Information

Gene ID	1139;89832
Other Names	CHRNA7; NACHRA7; Neuronal acetylcholine receptor subunit alpha-7; CHRFA7A; CHRNA7-FAM7A fusion protein; CHRNA7-DR1; D-10
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CHRNA7. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CHRNA7 (HGNC:1960)
Synonyms	NACHRA7
Function	Component of neuronal acetylcholine receptors (nAChRs) that function as pentameric, ligand-gated cation channels with high calcium permeability among other activities. nAChRs are excitatory neurotransmitter receptors formed by a collection of nAChR subunits known to mediate synaptic transmission in the nervous system and the neuromuscular junction. Each nAChR subunit confers differential attributes to channel properties, including activation, deactivation and desensitization kinetics, pH sensitivity, cation permeability, and binding to allosteric modulators (PubMed: 15609996 , PubMed: 33735609 , PubMed: 8145738). CHRNA7 forms homopentameric neuronal acetylcholine receptors abundantly expressed in the central nervous system, characterized by fast desensitization and high calcium permeability (PubMed: 31560909 , PubMed: 33735609 , PubMed: 38382524 ,

PubMed:[8145738](#)). Also forms heteropentamers with CHRNB2, mainly expressed in basal forebrain cholinergic neurons. Involved in the modulation of calcium- dependent signaling pathways and influences the release of neurotransmitters, including dopamine, glutamate and GABA (PubMed:[33239400](#)). Also expressed in non-neuronal cells such as immune cells like lymphocytes, monocytes and macrophages (PubMed:[12508119](#), PubMed:[16968406](#), PubMed:[25259522](#)). In T cells, activation induces metabotropic signaling that results in an increase of intracellular Ca²⁺ concentrations, independent of ionotropic receptor functions (PubMed:[17709503](#)). In macrophages, required for acetylcholine-mediated inhibition of TNF and other inflammatory cytokine release (PubMed:[12508119](#)). Once activated by acetylcholine, nicotine or other agonists, selectively inhibits production of pro-inflammatory cytokines while leaving anti-inflammatory cytokines undisturbed (PubMed:[12508119](#), PubMed:[25259522](#)). Stimulates the cholinergic anti-inflammatory pathway, controlling inflammation by inhibiting NFκB nuclear translocation and activating the JAK2-STAT3 pathway, independently of ion channel activity (PubMed:[16968406](#), PubMed:[25259522](#)). Also expressed in the urothelium where it modulates reflex bladder activity by increasing intracellular calcium through internal stores and decreasing basal ATP release (By similarity).


Cellular Location

Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q05941}; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=TMEM35A/NACHO promotes its trafficking to the cell membrane (PubMed:27789755). RIC3 promotes its trafficking to the cell membrane (By similarity) {ECO:0000250|UniProtKB:Q05941, ECO:0000269|PubMed:27789755}

Tissue Location

Expressed in neuronal cells (PubMed:8145738). Expressed in macrophages (at protein level) (PubMed:12508119)

Images



Western blot analysis of CHRNA7 expression in mouse liver (A), rat liver (B) whole cell lysates. (Predicted band size: 11; 56; 59 kD; Observed band size: 46 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.