

Anti-ENDOG Antibody

Catalog # AP53667

Product Information

Application WB Primary Accession Q14249

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW32620

Additional Information

Gene ID 2021

Other Names Endonuclease G, mitochondrial; Endo G

Target/Specificity Recognizes endogenous levels of ENDOG protein.

Dilution WB~~1/500 - 1/1000

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name ENDOG

Function Endonuclease that preferentially catalyzes the cleavage of double-stranded


5-hydroxymethylcytosine (5hmC)-modified DNA (PubMed:25355512). The 5hmC-modified nucleotide does not increase the binding affinity, but instead increases the efficiency of cutting and specifies the site of cleavage for the modified DNAs (By similarity). Shows significantly higher affinity for four-stranded Holliday junction over duplex and single-stranded DNAs (By similarity). Promotes conservative recombination when the DNA is 5hmC-modified (PubMed:25355512). Promotes autophagy through the suppression of mTOR by its phosphorylation-mediated interaction with YWHAG and its endonuclease activity-mediated DNA damage response (PubMed:33473107). GSK3-beta mediated phosphorylation of ENDOG enhances its interaction with YWHAG, leading to the release of TSC2 and PIK3C3 from YWHAG resulting in mTOR pathway suppression and autophagy initiation (PubMed:33473107). Promotes cleavage of mtDNA in response to oxidative and nitrosative stress, in turn inducing compensatory mtDNA replication (PubMed:29719607).

replication (Fubined.<u>297 19007)</u>

Background

Rabbit polyclonal antibody to ENDOG

Images



Western blot analysis of ENDOG expression in NCIH292 (A), HepG2 (B), Raw264.7 (C), PC12 (D) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.