

Anti-ARD1 Antibody

Catalog # AP53696

Product Information

Application	WB, IF/IC
Primary Accession	Q12972
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38479

Additional Information


Gene ID	5511
Other Names	ARD1; NIPP1; Nuclear inhibitor of protein phosphatase 1; NIPP-1; Protein phosphatase 1 regulatory inhibitor subunit 8
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ARD1. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	PPP1R8
Synonyms	ARD1, NIPP1
Function	Inhibitor subunit of the major nuclear protein phosphatase-1 (PP-1). It has RNA-binding activity but does not cleave RNA and may target PP-1 to RNA-associated substrates. May also be involved in pre- mRNA splicing. Binds DNA and might act as a transcriptional repressor. Seems to be required for cell proliferation.
Cellular Location	Nucleus. Nucleus speckle. Note=Primarily, but not exclusively, nuclear
Tissue Location	Ubiquitously expressed, with highest levels in heart and skeletal muscle, followed by brain, placenta, lung, liver and pancreas. Less abundant in kidney. The concentration and ratio between isoforms is cell-type dependent. Isoform Alpha (>90%) and isoform Beta were found in brain, heart and kidney. Isoform Gamma is mainly found in B-cells and T-lymphocytes, and has been

found in 293 embryonic kidney cells.

Images



Western blot analysis of ARD1 expression in HEK293T (A), MCF7 (B), mouse testis (C), rat testis (D) whole cell lysates. (Predicted band size: 38 kD; Observed band size: 41 kD)



Immunofluorescent analysis of ARD1 staining in PANC1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.