

Anti-NRSN2 Antibody

Catalog # AP53743

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q9GZP1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21983

Additional Information



Gene ID	80023
Other Names	C20orf98; Neurensin-2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NRSN2. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	NRSN2
Synonyms	C20orf98
Function	May play a role in maintenance and/or transport of vesicles.
Cellular Location	Membrane; Multi-pass membrane protein

Images

Western blot analysis of NRSN2 expression in HepG2 (A), HCT116 (B), AML12 (C), PC12 (D) whole cell lysates.
(Predicted band size: 21 kD; Observed band size: 34 kD)



Immunohistochemical analysis of NRSN2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of NRSN2 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.