

Anti-ZHX2 Antibody

Catalog # AP53783

Product Information

Application	WB
Primary Accession	Q9Y6X8
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92307

Additional Information

Gene ID	22882
Other Names	AFR1; KIAA0854; RAF; Zinc fingers and homeoboxes protein 2; Alpha-fetoprotein regulator 1; AFP regulator 1; Regulator of AFP; Zinc finger and homeodomain protein 2
Target/Specificity	Recognizes endogenous levels of ZHX2 protein.
Dilution	WB~~1/500 - 1/1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ZHX2
Synonyms	AFR1, KIAA0854, RAF
Function	Acts as a transcriptional repressor (PubMed: 12741956). Represses the promoter activity of the CDC25C gene stimulated by NFYA (PubMed: 12741956). May play a role in retinal development where it regulates the composition of bipolar cell populations, by promoting differentiation of bipolar OFF-type cells (By similarity). In the brain, may promote maintenance and suppress differentiation of neural progenitor cells in the developing cortex (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00108, ECO:0000269 PubMed:12741956, ECO:0000269 PubMed:17056598} Note=Colocalizes with EFNB1 intracellular domain in the nucleus {ECO:0000250 UniProtKB:Q8C0C0}


Tissue Location

Ubiquitously expressed. Expressed in podocytes.


Background

Rabbit polyclonal antibody to ZHX2

Images



Western blot analysis of ZHX2 expression in HepG2 (A), K562 (B) whole cell lysates.



Immunohistochemical analysis of ZHX2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.