

Anti-ADPGK Antibody

Catalog # AP53786

Product Information

ApplicationWB, IFPrimary AccessionQ9BRR6

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW54089

Additional Information

Gene ID 83440

Other Names ADP-dependent glucokinase; ADP-GK; ADPGK; RbBP-35

Target/Specificity Recognizes endogenous levels of ADPGK protein.

Dilution WB~~1/500 - 1/1000 IF~~1/50 - 1/200

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name ADPGK

Function Catalyzes the phosphorylation of D-glucose to D-glucose 6- phosphate using

ADP as the phosphate donor. GDP and CDP can replace ADP, but with

reduced efficiency (By similarity).


Cellular Location Secreted.


Background

Rabbit polyclonal antibody to ADPGK


Images

Western blot analysis of ADPGK expression in HeLa (A), VEC (B) whole cell lysates.





Immunohistochemical analysis of ADPGK staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ADPGK staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.