

# Anti-SIAH1/2 Antibody

Catalog # AP53797

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IF                 |
| Primary Accession | <a href="#">Q8IUQ4</a> |
| Other Accession   | <a href="#">Q43255</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 31123                  |

## Additional Information

---

|                    |  |
|--------------------|--|
| Gene ID            | 6477   |
| Other Names        | SIAH1; HUMSIAH; E3 ubiquitin-protein ligase SIAH1; Seven in absentia homolog 1; Siah-1; Siah-1a; SIAH2; E3 ubiquitin-protein ligase SIAH2; Seven in absentia homolog 2; Siah-2; hSiah2 |
| Target/Specificity | Recognizes endogenous levels of SIAH1/2 protein.   |
| Dilution           | WB~~1/500 - 1/1000 IF~~1/50 - 1/200  |
| Format             | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.  |
| Storage            | Store at -20 °C.Stable for 12 months from date of receipt  |

## Protein Information

---

|          |  |
|----------|--|
| Name     | SIAH1  |
| Synonyms | HUMSIAH  |
| Function | <p>E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:<a href="#">14506261</a>, PubMed:<a href="#">14645235</a>, PubMed:<a href="#">14654780</a>, PubMed:<a href="#">15064394</a>, PubMed:<a href="#">16085652</a>, PubMed:<a href="#">19224863</a>, PubMed:<a href="#">20508617</a>, PubMed:<a href="#">22483617</a>, PubMed:<a href="#">28546513</a>, PubMed:<a href="#">32430360</a>, PubMed:<a href="#">33591310</a>, PubMed:<a href="#">9334332</a>, PubMed:<a href="#">9858595</a>). E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:<a href="#">14506261</a>, PubMed:<a href="#">14645235</a>, PubMed:<a href="#">14654780</a>, PubMed:<a href="#">15064394</a>, PubMed:<a href="#">16085652</a>, PubMed:<a href="#">19224863</a>, PubMed:<a href="#">20508617</a>, PubMed:<a href="#">22483617</a>, PubMed:<a href="#">9334332</a>, PubMed:<a href="#">9858595</a>).</p> <p>Mediates E3 ubiquitin ligase activity either through direct binding to</p> |

substrates or by functioning as the essential RING domain subunit of larger E3 complexes (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (ELL2, MYB, POU2AF1, PML and RBBP8), a cell surface receptor (DCC), the cell-surface receptor-type tyrosine kinase FLT3, the cytoplasmic signal transduction molecules (KLF10/TIEG1 and NUMB), an antiapoptotic protein (BAG1), a microtubule motor protein (KIF22), a protein involved in synaptic vesicle function in neurons (SYP), a structural protein (CTNNB1) and SNCAIP (PubMed:[10747903](#), PubMed:[11146551](#), PubMed:[11389839](#), PubMed:[11389840](#), PubMed:[11483517](#), PubMed:[11483518](#), PubMed:[11752454](#), PubMed:[12072443](#)). Confers constitutive instability to HIPK2 through proteasomal degradation (PubMed:[18536714](#), PubMed:[33591310](#)). It is thereby involved in many cellular processes such as apoptosis, tumor suppression, cell cycle, axon guidance, transcription regulation, spermatogenesis and TNF-alpha signaling (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Has some overlapping function with SIAH2 (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Induces apoptosis in cooperation with PEG3 (By similarity). Upon nitric oxid (NO) generation that follows apoptotic stimulation, interacts with S-nitrosylated GAPDH, mediating the translocation of GAPDH to the nucleus (By similarity). GAPDH acts as a stabilizer of SIAH1, facilitating the degradation of nuclear proteins (By similarity). Mediates ubiquitination and degradation of EGLN2 and EGLN3 in response to the unfolded protein response (UPR), leading to their degradation and subsequent stabilization of ATF4 (By similarity). Also part of the Wnt signaling pathway in which it mediates the Wnt-induced ubiquitin- mediated proteasomal degradation of AXIN1 (PubMed:[28546513](#), PubMed:[32430360](#)).

#### Cellular Location

Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Partially nuclear


#### Tissue Location

Widely expressed at a low level. Down-regulated in advanced hepatocellular carcinomas.


## Background

Rabbit polyclonal antibody to SIAH1/2


## Images



Western blot analysis of SIAH1/2 expression in mouse muscle (A), rat muscle (B) whole cell lysates.



Immunohistochemical analysis of SIAH1/2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of SIAH1/2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.