

Anti-SP3/4 Antibody

Catalog # AP53798

Product Information

Application	WB, IF
Primary Accession	Q02447
Other Accession	Q02446
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	81925

Additional Information

Gene ID	6670
Other Names	SP3; Transcription factor Sp3; SPR-2; SP4; Transcription factor Sp4; SPR-1
Target/Specificity	Recognizes endogenous levels of SP3/4 protein.
Dilution	WB~~1/500 - 1/1000 IF~~1/50 - 1/200
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	SP3
Function	Transcriptional factor that can act as an activator or repressor depending on isoform and/or post-translational modifications. Binds to GT and GC boxes promoter elements. Competes with SP1 for the GC-box promoters. Weak activator of transcription but can activate a number of genes involved in different processes such as cell-cycle regulation, hormone-induction and house-keeping.
Cellular Location	Nucleus. Nucleus, PML body. Note=Localizes to the nuclear periphery and in nuclear dots when sumoylated. Some localization in PML nuclear bodies
Tissue Location	Ubiquitously expressed.


Background

Rabbit polyclonal antibody to SP3/4

Images



Western blot analysis of SP3/4 expression in HeLa (A), K562 (B) whole cell lysates.



Immunofluorescent analysis of SP3/4 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.