

Anti-MAP126 Antibody

Catalog # AP53823

Product Information

Application	WB, IF/IC
Primary Accession	Q96R06
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	134422

Additional Information

Gene ID	10615
Other Names	Sperm-associated antigen 5; Astrin; Deepest; Mitotic spindle-associated protein p126; MAP126
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human MAP126. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SPAG5
Function	Essential component of the mitotic spindle required for normal chromosome segregation and progression into anaphase (PubMed: 11724960 , PubMed: 12356910 , PubMed: 27462074). Required for chromosome alignment, normal timing of sister chromatid segregation, and maintenance of spindle pole architecture (PubMed: 17664331 , PubMed: 27462074). In complex with SKAP, promotes stable microtubule- kinetochore attachments. May contribute to the regulation of separase activity. May regulate AURKA localization to mitotic spindle, but not to centrosomes and CCNB1 localization to both mitotic spindle and centrosomes (PubMed: 18361916 , PubMed: 21402792). Involved in centriole duplication. Required for CDK5RAP2, CEP152, WDR62 and CEP63 centrosomal localization and promotes the centrosomal localization of CDK2 (PubMed: 26297806). In non-mitotic cells, upon stress induction, inhibits mammalian target of rapamycin complex 1 (mTORC1) association and recruits the mTORC1 component RPTOR to stress granules (SGs), thereby preventing mTORC1 hyperactivation-induced apoptosis

(PubMed:[23953116](#)). May enhance GSK3B-mediated phosphorylation of other substrates, such as MAPT/TAU (PubMed:[18055457](#)).


Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Chromosome, centromere, kinetochore. Midbody Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Cytoplasmic granule. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite
Note=Colocalizes with PCM1 at centriolar satellites throughout the cell cycle (PubMed:26297806). In a punctate pattern in interphase cells During mitosis, detected at spindle poles during prophase, throughout the spindle in metaphase and anaphase, and at midzone microtubules in anaphase and telophase (PubMed:27462074). Efficient targeting to the mitotic spindle may depend upon phosphorylation by GSK3B. Detected on kinetochores of chromosomes that have congressed. The astrin (SPAG5)- kinastrin (SKAP) complex localizes to the microtubule plus ends (By similarity). In non-mitotic non-stressed cells, shows a microtubuli pattern. In arsenite-stressed cells, accumulates in stress granules {ECO:0000250, ECO:0000269 | PubMed:26297806, ECO:0000269 | PubMed:27462074}


Tissue Location

Highly expressed in testis. Detected at low levels in placenta, liver, pancreas, thymus and colon

Images



Western blot analysis of MAP126 expression in Myla2059 (A), HEK293T (B), HuT78 (C) whole cell lysates. (Predicted band size: 134 kD; Observed band size: 135; 150 kD)



Immunofluorescent analysis of MAP126 staining in MDAMB231 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.