

Anti-RPL3L Antibody

Catalog # AP53831

Product Information

Application	WB, IF
Primary Accession	Q92901
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46296

Additional Information

Gene ID	6123
Other Names	60S ribosomal protein L3-like
Target/Specificity	Recognizes endogenous levels of RPL3L protein.
Dilution	WB~~1/500 - 1/1000 IF~~1/50 - 1/200
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	RPL3L {ECO:0000303 PubMed:8921388, ECO:0000312 HGNC:HGNC:10351}
Function	Heart- and skeletal muscle-specific component of the ribosome, which regulates muscle function. Component of the large ribosomal subunit in striated muscle cells: replaces the RPL3 paralog in the ribosome in these cells. The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell. Inhibits myotube growth and muscle function.

Background


Rabbit polyclonal antibody to RPL3L

Images


Western blot analysis of RPL3L expression in Jurkat (A),



Hela (B) whole cell lysates.



Immunohistochemical analysis of RPL3L staining in human muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of RPL3L staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.