

Anti-RNF138 Antibody

Catalog # AP53865

Product Information

Application	WB
Primary Accession	Q8WVD3
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28193

Additional Information

Gene ID	51444
Other Names	NARF; E3 ubiquitin-protein ligase RNF138; Nemo-like kinase-associated RING finger protein; NLK-associated RING finger protein; hNARF; RING finger protein 138
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human RNF138. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	RNF138 (HGNC:17765)
Function	E3 ubiquitin-protein ligase involved in DNA damage response by promoting DNA resection and homologous recombination (PubMed:26502055, PubMed:26502057). Recruited to sites of double-strand breaks following DNA damage and specifically promotes double-strand break repair via homologous recombination (PubMed:26502055, PubMed:26502057). Two different, non-exclusive, mechanisms have been proposed. According to a report, regulates the choice of double-strand break repair by favoring homologous recombination over non-homologous end joining (NHEJ): acts by mediating ubiquitination of XRCC5/Ku80, leading to remove the Ku complex from DNA breaks, thereby promoting homologous recombination (PubMed:26502055). According to another report, cooperates with UBE2Ds E2 ubiquitin ligases (UBE2D1, UBE2D2, UBE2D3 or UBE2D4) to promote homologous recombination by mediating ubiquitination of RBBP8/CtIP (PubMed:26502057). Together with NLK, involved in the ubiquitination and

degradation of TCF/LEF (PubMed:[16714285](#)). Also exhibits auto-ubiquitination activity in combination with UBE2K (PubMed:[16714285](#)). May act as a negative regulator in the Wnt/beta-catenin-mediated signaling pathway (PubMed:[16714285](#)).


Cellular Location

Chromosome. Note=Recruited at DNA damage sites (PubMed:26502055). Localizes to sites of double-strand break: localization to double-strand break sites is mediated by the zinc fingers (PubMed:26502055, PubMed:26502057)

Background

Rabbit polyclonal antibody to RNF138

Images



Western blot analysis of RNF138 expression in CT26 (A), PC12 (B), COS7 (C), MCF7 (D), A549 (E) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.