

# Anti-Collagen 18 alpha 1 Antibody

Catalog # AP53893

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IF                 |
| Primary Accession | <a href="#">P39060</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 178188                 |

## Additional Information

---

|                    |  |
|--------------------|--|
| Gene ID            | 80781  |
| Other Names        | Collagen alpha-1(XVIII) chain  |
| Target/Specificity | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Collagen 18 alpha 1. The exact sequence is proprietary. |
| Dilution           | WB~~1/500 - 1/1000 IF~~1/50 - 1/200  |
| Format             | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.                                    |
| Storage            | Store at -20 °C.Stable for 12 months from date of receipt  |

## Protein Information


---

|                   |   |
|-------------------|---|
| Name              | COL18A1 ( <a href="#">HGNC:2195</a> )   |
| Function          | Probably plays a major role in determining the retinal structure as well as in the closure of the neural tube.  |
| Cellular Location | Secreted, extracellular space, extracellular matrix. Secreted, extracellular space, extracellular matrix, basement membrane {ECO:0000250 UniProtKB:P39061} [Endostatin]: Secreted. Secreted, extracellular space, extracellular matrix, basement membrane |
| Tissue Location   | Detected in placenta (at protein level) (PubMed:32337544). Present in multiple organs with highest levels in liver, lung and kidney.  |


## Background

---


Rabbit polyclonal antibody to Collagen 18 alpha 1



Western blot analysis of Collagen 18 alpha 1 expression in MCF7 (A) whole cell lysates.



Immunohistochemical analysis of Collagen 18 alpha 1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Collagen 18 alpha 1 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

## Citations

---

- [Discovery of IgG4 Anti-Gliadin Autoantibody as a Potential Biomarker of Psoriasis Using an Autoantigen Array](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.