

Anti-ASF1B Antibody

Catalog # AP53915

Product Information

Application	WB, IHC
Primary Accession	Q9NVP2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22434

Additional Information

Gene ID	55723
Other Names	Histone chaperone ASF1B; Anti-silencing function protein 1 homolog B; hAsf1; hAsf1b; CCG1-interacting factor A-II; CIA-II; hCIA-II
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ASF1B. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IHC~~1/50 - 1/200
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ASF1B {ECO:0000303 PubMed:11470414, ECO:0000312 HGNC:HGNC:20996}
Function	Histone chaperone that facilitates histone deposition and histone exchange and removal during nucleosome assembly and disassembly (PubMed: 11897662 , PubMed: 14718166 , PubMed: 15664198 , PubMed: 16151251 , PubMed: 21454524 , PubMed: 26527279). Cooperates with chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly (PubMed: 11897662 , PubMed: 14718166 , PubMed: 15664198 , PubMed: 16151251). Also involved in the nuclear import of the histone H3-H4 dimer together with importin-4 (IPO4): specifically recognizes and binds newly synthesized histones with the monomethylation of H3 'Lys-9' (H3K9me1) and diacetylation at 'Lys-5' and 'Lys-12' of H4 (H4K5K12ac) marks in the cytosol (PubMed: 20953179 , PubMed: 21454524 , PubMed: 26527279). Does not participate in replication-independent nucleosome deposition which is mediated by ASF1A and HIRA (PubMed: 11897662 , PubMed: 14718166 , PubMed: 15664198 , PubMed: 16151251). Required for gonad development (PubMed: 12842904).


Cellular Location

Nucleus. Cytoplasm, cytosol


Tissue Location

Highly expressed in testis and at lower levels in colon, small intestine and thymus.


Images



Western blot analysis of ASF1B expression in HEK293T (A), U87MG (B), Hela (C), K562 (D), MCF7 (E) whole cell lysates. (Predicted band size: 22 kD; Observed band size: 22 kD)



Immunohistochemical analysis of ASF1B staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of ASF1B staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.