

Anti-MCT12 Antibody

Catalog # AP53919

Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q6ZSM3 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 56498 |

Additional Information


| | |
|--------------------|---|
| Gene ID | 387700 |
| Other Names | MCT12; Monocarboxylate transporter 12; MCT 12; Solute carrier family 16 member 12 |
| Target/Specificity | Recognizes endogenous levels of MCT12 protein. |
| Dilution | WB~~1/500 - 1/1000 |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |

Protein Information


| | |
|-------------------|---|
| Name | SLC16A12 (HGNC:23094) |
| Function | Functions as a transporter for creatine and as well for its precursor guanidinoacetate. Transport of creatine and GAA is independent of resting membrane potential and extracellular Na(+), Cl(-), or pH. Contributes to the process of creatine biosynthesis and distribution. |
| Cellular Location | Cell membrane; Multi-pass membrane protein. Basolateral cell membrane {ECO:0000250 UniProtKB:Q8BGC3}; Multi-pass membrane protein. Note=Interaction with isoform 2 of BSG is required for its localization to the plasma membrane. |
| Tissue Location | Most highly expressed in kidney, followed by retina, lung, heart and testis. Very weakly expressed in brain and liver. Also detected in lens. |

Background

Images



Western blot analysis of MCT12 expression in Jurkat (A) whole cell lysates.



Immunohistochemical analysis of MCT12 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.