

Anti-TBL2 Antibody

Catalog # AP53923

Product Information

Application	WB, IHC
Primary Accession	Q9Y4P3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49798

Additional Information



Gene ID	26608
Other Names	WBSCR13; Transducin beta-like protein 2; WS beta-transducin repeats protein; WS-betaTRP; Williams-Beuren syndrome chromosomal region 13 protein
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TBL2. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IHC~~1/50 - 1/200
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	TBL2
Synonyms	WBSCR13

Images

Western blot analysis of TBL2 expression in HEK293T (A), U87MG (B), MCF7 (C) whole cell lysates. (Predicted band size: 49 kD; Observed band size: 50 kD)



Immunohistochemical analysis of TBL2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of TBL2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.