

# Anti-MARCH2 Antibody

Catalog # AP53932

## Product Information

Application	WB, IF
Primary Accession	<a href="#">Q9P0N8</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26995

## Additional Information

Gene ID	51257
Other Names	RNF172; E3 ubiquitin-protein ligase MARCH2; Membrane-associated RING finger protein 2; Membrane-associated RING-CH protein II; MARCH-II; RING finger protein 172
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MARCH2. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/2000 IF~~1/50 - 1/100
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	MARCHF2 ( <a href="#">HGNC:28038</a> )
Synonyms	MARCH2, RNF172
Function	E3 ubiquitin-protein ligase that may mediate ubiquitination of TFRC and CD86, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates (PubMed: <a href="#">14722266</a> , PubMed: <a href="#">16428329</a> ). Together with GOPC/CAL mediates the ubiquitination and lysosomal degradation of CFTR (PubMed: <a href="#">23818989</a> ). Ubiquitinates and therefore mediates the degradation of DLG1 (PubMed: <a href="#">17980554</a> ). Regulates the intracellular trafficking and secretion of alpha1-antitrypsin/SERPINA1 and HP/haptoglobin via ubiquitination and degradation of the cargo receptor ERGIC3 (PubMed: <a href="#">31142615</a> ). Negatively regulates the antiviral and antibacterial immune response by repression of the NF-kB and type 1 IFN

signaling pathways, via MARCHF2-mediated K48-linked polyubiquitination of IKBKG/NEMO, resulting in its proteasomal degradation (PubMed:[32935379](#)). May be involved in endosomal trafficking through interaction with STX6 (PubMed:[15689499](#)).

### Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Lysosome membrane; Multi-pass membrane protein. Endosome membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasm. Cell membrane; Multi-pass membrane protein


### Tissue Location

Broadly expressed..


## Background

Rabbit polyclonal antibody to MARCH2


## Images



Western blot analysis of MARCH2 expression in mouse brain (A), rat spleen (B), mouse kidney (C), rat kidney (D) whole cell lysates.



Immunohistochemical analysis of MARCH2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of MARCH2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.