

Anti-TAGAP Antibody

Catalog # AP53934

Product Information

| | |
|--------------------------|------------------------|
| Application | WB |
| Primary Accession | Q8N103 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 80703 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 117289 |
| Other Names | TAGAP1; T-cell activation Rho GTPase-activating protein; T-cell activation GTPase-activating protein |
| Target/Specificity | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TAGAP. The exact sequence is proprietary. |
| Dilution | WB~~1/500 - 1/2000 |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |

Protein Information


| | |
|-----------------|--|
| Name | TAGAP |
| Synonyms | TAGAP1 |
| Function | May function as a GTPase-activating protein and may play important roles during T-cell activation. |

Background


Rabbit polyclonal antibody to TAGAP

Images

Western blot analysis of TAGAP expression in mouse



kidney (A), rat kidney (B), rat liver (C), rat heart (D) whole cell lysates.



Immunohistochemical analysis of TAGAP staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.