

Anti-UEVLD Antibody

Catalog # AP53985

Product Information

Application	WB, IHC
Primary Accession	Q8IX04
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52264

Additional Information



Gene ID	55293
Other Names	UEV3; Ubiquitin-conjugating enzyme E2 variant 3; UEV-3; EV and lactate/malate dehydrogenase domain-containing protein
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human UEVLD. The exact sequence is proprietary.
Dilution	WB~~1/500 - 1/1000 IHC~~1/50 - 1/100
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	UEVLD {ECO:0000312 EMBL:AAH64566.1}
Function	Possible negative regulator of polyubiquitination.
Tissue Location	Colon, colon carcinoma cell lines, normal cervical epithelium, carcinomas of the uterine cervix and peripheral blood leukocytes.

Images

Western blot analysis of UEVLD expression in Hela (A), Jurkat (B), H9C2 (C), AML12 (D) whole cell lysates.
(Predicted band size: 52 kD; Observed band size: 52 kD)



Immunohistochemical analysis of UEVLD staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of UEVLD staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.