

# Anti-GITRL Antibody

Catalog # AP54053

#### **Product Information**

Application WB
Primary Accession Q9UNG2

**Reactivity** Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 20308

#### **Additional Information**

**Gene ID** 8995

Other Names AITRL; GITRL; TL6; Tumor necrosis factor ligand superfamily member 18;

Activation-inducible TNF-related ligand; AITRL; Glucocorticoid-induced

TNF-related ligand; hGITRL

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human GITRL. The exact sequence is proprietary.

**Dilution** WB~~1/500 - 1/1000

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name TNFSF18 ( HGNC:11932)

Synonyms AITRL, GITRL, TL6

**Function** Cytokine that binds to TNFRSF18/AITR/GITR. Regulates T-cell responses. Can

function as costimulator and lower the threshold for T- cell activation and

T-cell proliferation. Important for interactions between activated

T-lymphocytes and endothelial cells. Mediates activation of NF-kappa-B. Triggers increased phosphorylation of STAT1 and up-regulates expression of VCAM1 and ICAM1 (PubMed:23892569). Promotes leukocyte adhesion to endothelial cells (PubMed:23892569). Regulates migration of monocytes from

the splenic reservoir to sites of inflammation (By similarity).


**Cellular Location** Cell membrane; Single-pass type II membrane protein

**Tissue Location** Expressed at high levels in the small intestine, ovary, testis, kidney and

## **Background**

Rabbit polyclonal antibody to GITRL

### **Images**



Western blot analysis of GITRL expression in SKOVCAR3 (A), A2780 (B), HEK293T (C), mouse testis (D), rat testis (E) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.