

Anti-MXD4 Antibody

Catalog # AP54092

Product Information

Application WB
Primary Accession Q14582

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW23528

Additional Information

Gene ID 10608

Other Names BHLHC12; MAD4; Max dimerization protein 4; Max dimerizer 4; Class C basic

helix-loop-helix protein 12; bHLHc12; Max-associated protein 4;

Max-interacting transcriptional repressor MAD4

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the

N-term region of human MXD4. The exact sequence is proprietary.

Dilution WB~~1/500 - 1/1000

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name MXD4

Synonyms BHLHC12, MAD4

Function Transcriptional repressor. Binds with MAX to form a sequence- specific


DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. Antagonizes MYC transcriptional activity by competing for MAX and suppresses MYC dependent cell transformation (By similarity).

Cellular Location Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00981}.


Background

Rabbit polyclonal antibody to MXD4

Images



Western blot analysis of MXD4 expression in NIH3T3 (A), PMVEC (B), Hela (C), A375 (D) whole cell lysates.



Immunohistochemical analysis of MXD4 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.