10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CENPA (Phospho-S7) Antibody

Catalog # AP54111

Product Information

Application	WB, IF/IC
Primary Accession	P49450
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15991

Additional Information

Gene ID	1058
Other Names	Histone H3-like centromeric protein A; Centromere autoantigen A; Centromere protein A; CENP-A
Target/Specificity	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding S7 of human CENPA protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CENPA
Function	Histone H3-like nucleosomal protein that is specifically found in centromeric nucleosomes (PubMed:11756469, PubMed:14667408, PubMed:15282608, PubMed:15475964, PubMed:15702419, PubMed:17651496, PubMed:19114591, PubMed:20739937, PubMed:27499292, PubMed:7962047, PubMed:9024683). Replaces conventional H3 in the nucleosome core of centromeric chromatin that serves as an assembly site for the inner kinetochore (PubMed:18072184). The presence of CENPA subtly modifies the nucleosome structure and the way DNA is wrapped around the nucleosome and gives rise to protruding DNA ends that are less well- ordered and rigid compared to nucleosomes containing histone H3 (PubMed:26878239, PubMed:27499292). May serve as an epigenetic mark that propagates centromere identity through replication and cell division (PubMed:15282608, PubMed:15475964, PubMed:20739937, PubMed:21478274, PubMed:26878239). Required for recruitment and assembly of kinetochore proteins, and as a consequence required for progress through mitosis,

chromosome segregation and cytokinesis (PubMed:11756469, PubMed:14667408, PubMed:18072184, PubMed:23818633, PubMed:25556658, PubMed:27499292).


Cellular Location

Nucleus. Chromosome, centromere. Note=Localizes exclusively to sites of kinetochore assembly in centromeres. Occupies a compact domain at the inner kinetochore plate stretching across 2 thirds of the length of the constriction but encompassing only one third of the constriction width and height (PubMed:19114591) Phosphorylation at Ser-68 during early mitosis abolishes association with chromatin and centromeres and results in dispersed nuclear location (PubMed:25556658).


Images



Western blot analysis of CENPA (Phospho-S7) expression in HEK293T (A), Hela (B), DLD (C), rat testis (D), rat spleen (E) whole cell lysates. (Predicted band size: 15 kD; Observed band size: 17 kD)



Immunofluorescent analysis of CENPA (Phospho-S7) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Direct ELISA antibody dose-response curve using Anti-CENPA (Phospho-S7) Antibody. Antigen (Phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.