

Anti-OCT1 Antibody

Catalog # AP54135

Product Information

Application	WB
Primary Accession	P14859
Other Accession	P09086
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	76472

Additional Information


Gene ID	5451
Other Names	OCT1; OTF1; POU domain class 2 transcription factor 1; NF-A1; Octamer-binding protein 1; Oct-1; Octamer-binding transcription factor 1; OTF-1
Target/Specificity	Recognizes endogenous levels of OCT1 protein.
Dilution	WB~~1/500 - 1/2000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	POU2F1
Synonyms	OCT1, OTF1
Function	Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR.
Cellular Location	Nucleus.
Tissue Location	Ubiquitous. Isoform 2 is lymphocyte-specific.

Background

Images



Western blot analysis of 42278 expression in HeLa (A), Jurkat (B), HepG2 (C) whole cell lysates.



Immunohistochemical analysis of 42278 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.