10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NAIF1 Rabbit pAb

NAIF1 Rabbit pAb
Catalog # AP54198

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q69YI7
Reactivity	Human
Predicted	Rat, Dog, Mouse, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35164
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NAIF1
Epitope Specificity	3-85/327
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	NAIF protein 1 (nuclear apoptosis inducing factor 1; CB12-327; RP-379C10.2) belongs to the NAIF1 family. Apoptosis is a genetically determined cell suicidal pro gram that plays critical roles in many physiological and pathological processes. nuclear apoptosis-inducing factor 1 (NAIF1), overexpression of which induces apoptosis in cells. Human NAIF1 is located on chromosome 9q34.11 and encodes 327 amino acids with a homeodomain-like region and two nuclear localization signals at its N-terminal region. NAIF1 is conserved across diverse species, including human, mouse, crab-eating macaque, dog, chicken and frog, and shares no obvious homology to any known genes or proteins. Northern blot analysis revealed wide expression of NAIF1 mRNA throughout human tissues. NAIF1 was predominantly localized in the nucleus. Overexpression of NAIF1 inhibited cell growth and induced apoptosis. Furthermore, NAIF1 transfection caused both decreases in mitochondrial membrane potential and caspase-3 activation. In summary, NAIF1 is a nuclear protein that induces apoptosis when overexpressed.

Additional Information

Gene ID	203245
Other Names	Nuclear apoptosis-inducing factor 1, NAIF1, C9orf90
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	NAIF1
Synonyms	C9orf90
Function	Induces apoptosis.
Cellular Location	Nucleus
Tissue Location	Widely expressed..

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Tissue/cell: human colon tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37℃ for 20 min; Incubation: Anti-NAIF1 Polyclonal Antibody, Unconjugated(AP54198) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.