

# SLC33A1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54204

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, IF, E
<b>Primary Accession</b>	<a href="#">O00400</a>
<b>Reactivity</b>	Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	60909
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human SLC33A1
<b>Epitope Specificity</b>	481-549/549
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Endoplasmic reticulum membrane; Multi-pass membrane protein (Probable).
<b>SIMILARITY</b>	Belongs to the SLC33A transporter family.
<b>DISEASE</b>	Defects in SLC33A1 are the cause of spastic paraplegia autosomal dominant type 42 (SPG42) [MIM:612539]. Spastic paraplegia is a neurodegenerative disorder characterized by a slow, gradual, progressive weakness and spasticity of the lower limbs. Rate of progression and the severity of symptoms are quite variable. Initial symptoms may include difficulty with balance, weakness and stiffness in the legs, muscle spasms, and dragging the toes when walking. In some forms of the disorder, bladder symptoms (such as incontinence) may appear, or the weakness and stiffness may spread to other parts of the body
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	Acetyl-coenzyme A transporter 1 is required for the formation of O-acetylated (Ac) gangliosides. It is predicted to contain 6 to 10 transmembrane domains, and a leucine zipper motif in transmembrane domain III. Studies indicate that the protein is localized to the cytoplasm.

## Additional Information

---

<b>Gene ID</b>	9197
<b>Other Names</b>	Acetyl-coenzyme A transporter 1, AT-1, Acetyl-CoA transporter 1, Solute carrier family 33 member 1, SLC33A1, ACATN, AT1
<b>Target/Specificity</b>	Ubiquitous. Detected in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. With strongest signals in pancreas.
<b>Dilution</b>	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000


-10000

<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

<b>Name</b>	SLC33A1 ( <a href="#">HGNC:95</a> )
<b>Synonyms</b>	ACATN, AT1
<b>Function</b>	Acetyl-CoA transporter that mediates active acetyl-CoA import through the endoplasmic reticulum (ER) membrane into the ER lumen where specific ER-based acetyl-CoA:lysine acetyltransferases are responsible for the acetylation of ER-based protein substrates, such as BACE1 (PubMed: <a href="#">20826464</a> , PubMed: <a href="#">24828632</a> ). Necessary for O-acetylation of gangliosides (PubMed: <a href="#">9096318</a> ).
<b>Cellular Location</b>	Endoplasmic reticulum membrane; Multi-pass membrane protein
<b>Tissue Location</b>	Ubiquitous. Detected in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. With strongest signals in pancreas.

## Images



Protein: heart(mouse) lysates at 30ug;  
Primary: rabbit Anti-SLC33A1 (bs-0669R) at 1:300;  
Secondary: HRP conjugated Goat-Anti-rabbit  
IgG(bs-0295G-HRP) at 1: 5000;  
Predicted band size:61 kD  
Observed band size:61 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.