

Thyroid peroxidase Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54256

Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Physical State Immunogen Epitope Specificity Isotype Purity	WB, IHC-P, IHC-F, IF, E P07202 Rat Rabbit Polyclonal 102963 Liquid KLH conjugated synthetic peptide derived from human Thyroid peroxidase 111-210/933 IgG affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein. Isoform 3: Cell surface.
SIMILARITY	Belongs to the peroxidase family. XPO subfamily. Contains 1 EGF-like domain.
SUBUNIT	Iodination and coupling of the hormonogenic tyrosines in thyroglobulin to
	yield the thyroid hormones T(3) and T(4).
Post-translational	Glycosylated. Heme is covalently bound through a H(2)O(2)-dependent
modifications	autocatalytic process. Heme insertion is important for the delivery of protein at the cell surface. Cleaved in its N-terminal part.
DISEASE	Note=An alternative splicing in the thyroperoxidase mRNA can cause Graves' disease. Defects in TPO are the cause of thyroid dyshormonogenesis 2A (TDH2A) [MIM:274500]. A disorder due to defective conversion of accumulated iodide to organically bound iodine. The iodide organification defect can be partial or complete.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a membrane-bound glycoprotein. The encoded protein acts as an enzyme and plays a central role in thyroid gland function. The protein functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Multiple transcript variants encoding distinct isoforms have been identified for this gene, but the full-length nature of some variants has not been determined. [provided by RefSeq, May 2011].

Additional Information

Other Names	Thyroid peroxidase, TPO, 1.11.1.8, TPO (<u>HGNC:12015</u>)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000 -10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TPO (<u>HGNC:12015</u>)
Function	Iodination and coupling of the hormonogenic tyrosines in thyroglobulin to yield the thyroid hormones T(3) and T(4).
Cellular Location	Membrane; Single-pass type I membrane protein.

Images



Tissue/cell: Rat brain; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Thyroid peroxidase Polyclonal Antibody, Unconjugated(AP54256) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample:

Lane 1: MCF-7 (Human) Cell Lysate at 30 ug Lane 2: K562 (Human) Cell Lysate at 30 ug Lane 3: A673 (Human) Cell Lysate at 30 ug Lane 4: A431 (Human) Cell Lysate at 30 ug Primary: Anti- Thyroid peroxidase (AP54256) at 1/1000 dilution Anti-beta-Actin (bs-0061R) at 1/2000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 105-110 kD Observed band size: 105 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.