

IL-9 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54262

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	P15248
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15909
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human IL-9
Epitope Specificity	51-144/144
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the IL-7/IL-9 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	3578
Other Names	Interleukin-9, IL-9, Cytokine P40, T-cell growth factor P40, IL9
Dilution	IHC-P=1:100-500, IHC-F=1:100-500, ICC=1:100-500, IF=1:100-500, ELISA=1:5000-10000
Format	0.01M TBS (pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name IL9

Function Multifunctional cytokine secreted mainly by T-helper 2 lymphocytes and also mast cells or NKT cells that plays important roles in the immune response against parasites (PubMed:[29742432](#)). Affects intestinal epithelial permeability and adaptive immunity (PubMed:[29742432](#)). In addition, induces the differentiation of specific T-cell subsets such as IL-17 producing helper T-cells (TH17) and also proliferation and differentiation of mast cells. Mechanistically, exerts its biological effects through a receptor composed of IL9R subunit and a signal transducing subunit IL2RG. Receptor stimulation results in the rapid activation of JAK1 and JAK3 kinase activities leading to STAT1, STAT3 and STAT5-mediated transcriptional programs. Induction of differentiation genes seems to be mediated by STAT1 alone, while protection of cells from apoptosis depends on STAT3 and STAT5.

Cellular Location Secreted.

Images



Paraformaldehyde-fixed, paraffin embedded (Human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IL-9) Polyclonal Antibody, Unconjugated (AP54262) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.