

Angiopoietin 4 Rabbit pAb

Angiopoietin 4 Rabbit pAb
Catalog # AP54315

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9Y264
Reactivity	Mouse
Predicted	Human, Rat, Chicken, Dog, Pig, Horse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56849
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ANGPT4
Epitope Specificity	410-503/503
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted (Probable).
SIMILARITY	Contains 1 fibrinogen C-terminal domain.
SUBUNIT	Homodimer; disulfide-linked. Interacts with TEK/TIE2.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Angiopoietin 4 is a glycosylated protein with a fibrinogen C terminal domain. It is induced under hypoxic conditions in endothelial cells and is the target of peroxisome proliferation activators. Angiopoietin 4 is a serum hormone directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity and also acts as an apoptosis survival factor for vascular endothelial cells. It may play a role in several cancers and it also has been shown to prevent the metastatic process by inhibiting vascular activity as well as tumor cell motility and invasiveness. Decreased expression of this protein has been associated with type 2 diabetes. Alternatively spliced transcript variants encoding different isoforms have been described

Additional Information

Gene ID	51378
Other Names	Angiopoietin-4, ANG-4, Angiopoietin-3, ANG-3, ANGPT4, ANG3, ANG4
Target/Specificity	Highly expressed in the lung with much lower levels found in other tissues.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ANGPT4
Synonyms	ANG3, ANG4
Function	Binds to TEK/TIE2, modulating ANGPT1 signaling. Can induce tyrosine phosphorylation of TEK/TIE2. Promotes endothelial cell survival, migration and angiogenesis.
Cellular Location	Secreted.
Tissue Location	Highly expressed in the lung with much lower levels found in other tissues

Background

Angiopoietin 4 is a glycosylated protein with a fibrinogen C terminal domain. It is induced under hypoxic conditions in endothelial cells and is the target of peroxisome proliferation activators. Angiopoietin 4 is a serum hormone directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity and also acts as an apoptosis survival factor for vascular endothelial cells. It may play a role in several cancers and it also has been shown to prevent the metastatic process by inhibiting vascular activity as well as tumor cell motility and invasiveness. Decreased expression of this protein has been associated with type 2 diabetes. Alternatively spliced transcript variants encoding different isoforms have been described

Images



Tissue/cell: mouse tumor tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal
goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-ANGPT3/ANG3/ANGPT4/ANG4 Polyclonal
Antibody, Unconjugated(AP54315) 1:200, overnight at
4°C, followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.