

Myosin-13 Rabbit pAb

Myosin-13 Rabbit pAb

Catalog # AP54320

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UKX3.2
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Rabbit, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Myosin-13
Epitope Specificity	1801-1900/1938
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, myofibril. Note=Thick filaments of the myofibrils.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information


Other Names	Myosin-13, Myosin heavy chain 13, Myosin heavy chain, skeletal muscle, extraocular, MyHC-EO, Myosin heavy chain, skeletal muscle, laryngeal, MyHC-IIL, Superfast myosin, MYH13
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Myosin-13) Polyclonal Antibody, Unconjugated (AP54320) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.