

PCDHA10/CNRN8 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54388

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9Y5I2
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102875
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PCDHA10/CNRN8
Epitope Specificity	301-400/948
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Single-pass type I membrane protein. Isoform 2: Secreted.
SIMILARITY	Contains 6 cadherin domains.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. These genes generate thousands of different protocadherin proteins through alternative promoter usage and cis-alternative splicing. PCDHA and PCDHG family members form oligomers, which also increases the diversity of PCDH proteins at the cell surface. All three gene clusters, PCDHA, PCDHB, and PCDHG show upregulated expression during brain development, and PCDHA is subsequently downregulated by myelination. Expression of all three clusters continues in the olfactory bulb, hippocampus and cerebellum until adulthood. Members of the PCDH family are potential targets in schizophrenia and bipolar disorder pathogenesis. PCDHA10 produces at least three isoforms by alternative splicing.

Additional Information

Gene ID	56139
Other Names	Protocadherin alpha-10, PCDH-alpha-10, PCDHA10, CNRS8
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name PCDHA10

Synonyms CNRS8

Function Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Cellular Location Cell membrane; Single-pass type I membrane protein

Images

**Sample:**

Hippocampus (Mouse) Lysate at 40 ug

Primary: Anti- PCDHA10/CNRS8 (AP54388) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 100 kD

Observed band size: 115 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.