

## PCDHGC3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54407

## **Product Information**

**Application** WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat
Host
Clonality
Polyclonal
Calculated MW
Physical State

Q9UN70
Rat
Rabbit
Polyclonal
101077
Liquid

Immunogen KLH conjugated synthetic peptide derived from human PCDHGC3/PCDH2

Epitope Specificity 251-350/934

Isotype IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Cell membrane. Single-pass type I membrane protein.

**SIMILARITY** Contains 6 cadherin domains.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Protocadherins are a large family of cadherin-like cell adhesion proteins that

are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDH2 (protocadherin-2), also known as PCDHGC3 (protocadherin gamma subfamily C, 3) or PC43, is a 934 amino acid single-pass type I membrane protein that contains six cadherin domains and belongs to the protocadherin gamma family. Functioning as a calcium-dependent cell-adhesion protein, PCDH2 is thought to be involved in the establishment and maintenance of neuronal connections within the brain. Multiple isoforms of PCDH2 exist due

to alternative splicing events.

## **Additional Information**

**Gene ID** 5098

Other Names Protocadherin gamma-C3, PCDH-gamma-C3, Protocadherin-2,

Protocadherin-43, PC-43, PCDHGC3, PCDH2

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

## **Protein Information**

Name PCDHGC3

Synonyms PCDH2

**Function** Potential calcium-dependent cell-adhesion protein. May be involved in the

establishment and maintenance of specific neuronal connections in the brain.

**Cellular Location** Cell membrane; Single-pass type I membrane protein

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.