

Syntaxin 13 Rabbit pAb

Syntaxin 13 Rabbit pAb

Catalog # AP54476

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from rat Syntaxin 13
Epitope Specificity	101-200/274
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane. Golgi apparatus > trans-Golgi network membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Syntaxins, a family of proteins involved in the fusion of synaptic vesicles with the plasma membrane, display broad tissue distribution and contain C-terminal hydrophobic domains that direct them to their respective intracellular compartments. Syntaxin 12, also known as STX12, STX13 or STX14, is a 276 amino acid single-pass membrane protein that contains one t-SNARE coiled-coil homology domain and belongs to the Syntaxin family. Syntaxin 12 regulates protein transport between late endosomes and the trans-Golgi network and interacts with ABC1 (ATP-binding cassette transporter A1), a protein that facilitates cellular release of choline-phospholipids and cholesterol to apoA-I (apolipoprotein A-I). Found in recycling endosomes, Syntaxin 12 colocalizes with transferrin receptor and is encoded by a gene located on human chromosome 1.


Additional Information

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Syntaxin 13) Polyclonal Antibody, Unconjugated (AP54476) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.