10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

A4GNT Rabbit pAb

A4GNT Rabbit pAb
Catalog # AP54478

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9UNA3
Predicted	Rat, Pig, Dog, Human, Mouse, Chicken, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39497
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human A4GNT
Epitope Specificity	131-180/340
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Golgi apparatus membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Alpha 1,4-N-acetylflucosaminyltransferase (Alpha4Gn-T) mediates the biosynthesis of mucin type glycoprotein (O-glycan). Alpha4Gn-T acts as the key enzyme for the formation of the unique glycan GlcNAcalpha1-4Galbeta1-R, and most efficiently transfers N-acetylglucosamine (GlcNAc) to core 2 branched O-glycans. Alpha4Gn-T is a single-pass type II membrane protein associated with the Golgi apparatus and contains the conserved DXD motif involved in catalytic activity. It is expressed in stomach and pancreas, as well as in gastric cancer cells. Alpha4Gn-T is not expressed in peripheral blood cells, making it a useful biomarker for pancreatic cancer. Alpha4Gn-T and Mucin 6 expression is upregulated in the gastric mucosa of H.pylori infected patients, which suggest the involvement of ?Gn-T in defense against H. pylori infection.

Additional Information

Gene ID	51146
Other Names	Alpha-1, 4-N-acetylglucosaminyltransferase, Alpha4GnT, 2.4.1.-, A4GNT (HGNC:17968)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	A4GNT (HGNC:17968)
Function	Catalyzes the transfer of N-acetylglucosamine (GlcNAc) residues from UDP-N-acetyl-alpha-D-glucosamine donnors to beta (1->4)- linked or beta-(1->3)-linked galactose residues on O-glycans like core 2 branched O-glycans (PubMed:10430883). Necessary for the synthesis of type III mucin which is specifically produced in the stomach, duodenum, and pancreatic duct (PubMed:10430883). May protect against inflammation-associated gastric adenocarcinomas (By similarity).
Cellular Location	Golgi apparatus membrane; Single- pass type II membrane protein
Tissue Location	Detected in stomach and pancreas.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.