

MC4 Receptor Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54493

Product Information

Application	WB, IHC-F, IF, ICC, E
Primary Accession	P32245
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36943
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MC4 Receptor
Epitope Specificity	1-50/332
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Multi-pass membrane protein.
SIMILARITY	Belongs to the G-protein coupled receptor 1 family.
SUBUNIT	Interacts with ATRNL1. Interacts with MGRN1, but does not undergo MGRN1-mediated ubiquitination; this interaction competes with GNAS-binding and thus inhibits agonist-induced cAMP production.
DISEASE	Defects in MC4R are a cause of obesity (OBESITY) [MIM:601665]. It is a condition characterized by an increase of body weight beyond the limitation of skeletal and physical requirements, as the result of excessive accumulation of body fat.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The melanocortin family comprises the Alpha-, Beta- and Gamma- melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane G protein-coupled proteins that activate adenylyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.

Additional Information


Gene ID	4160
Other Names	Melanocortin receptor 4, MC4-R, MC4R
Target/Specificity	Brain, placental, and gut tissues.

Dilution	WB=1:500-2000,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information



Name	MC4R
Function	Hormone receptor that acts as a key component of the leptin- melanocortin pathway at the intersection of homeostatic maintenance of energetic state (PubMed: 32327598 , PubMed: 33858992). Plays a role in regulating food intake: activation by a stimulating hormone such as anorexigenic alpha-melanocyte stimulating hormone (alpha-MSH) inhibits appetite, whereas binding to a natural antagonist like Agouti-related protein/AGRP promotes appetite. G-protein-coupled receptor that activates conventional G-proteins signaling leading to induction of anorexigenic signaling in the hypothalamus to result in negative energy balance (PubMed: 33858992). Regulates the firing activity of neurons from the hypothalamus by alpha-MSH and AGRP independently of G-proteins signaling by ligand-induced coupling of closure of inwardly rectifying potassium channel KCNJ13 (By similarity). In intestinal epithelial cells, plays a role in the inhibition of hepatic glucose production via nesfatin-1/NUCB2 leading to increased cyclic adenosine monophosphate (cAMP) levels and glucagon-like peptide 1 (GLP-1) secretion in the intestinal epithelium (PubMed: 39562740).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Brain, placental, and gut tissues.

Images



Sample:
 Embryo (Mouse) Lysate at 40 ug
 Primary: Anti- MC4 Receptor (AP54493) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 37 kD
 Observed band size: 52 kD


Sample: MCF-7 Cell (Human) Lysate at 40 ug
 Primary: Anti-MC4 Receptor (AP54493) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 37 kD
 Observed band size: 40 kD



Key	Name	Parameter	Gate
	RSC96-blank.028	FL1-H	G1
	bs-0295G-FITC(B)-RSC96-6.053	FL1-H	G1
	bs-0295P-FITC(B)-RSC96-6.054	FL1-H	G1
	bs-11417R-FITC(B)-RSC96-6.058	FL1-H	G1

Positive control: RSC96


Isotype Control Antibody: Rabbit IgG ; Secondary Antibody: Goat anti-rabbit IgG-FITC, Dilution: 1:100 in 1 X PBS containing 0.5% BSA ; Primary Antibody Dilution: 6 µg in 100 µL 1X PBS containing 0.5% BSA.



Blank control: 293T(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice).

Primary Antibody: Rabbit Anti-MC4 Receptor /AF488 Conjugated antibody (AP54493-AF488), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG/AF488(orange), used under the same conditions.



Blank control: 293T(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice). Primary Antibody: Rabbit Anti-MC4 Receptor /AF488 Conjugated antibody (AP54493-AF488), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG/AF488(orange), used under the same conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.