10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Adrenodoxin Rabbit pAb

Adrenodoxin Rabbit pAb
Catalog # AP54496

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P10109
Reactivity	Rat
Predicted	Human, Mouse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19393
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Adrenodoxin/FDX
Epitope Specificity	51-150/184
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion matrix.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a small iron-sulfur protein that transfers electrons from NADPH through ferredoxin reductase to mitochondrial cytochrome P450, involved in steroid, vitamin D, and bile acid metabolism. Pseudogenes of this functional gene are found on chromosomes 20 and 21. [provided by RefSeq, Aug 2011]

Additional Information

Gene ID	2230
Other Names	Adrenodoxin, mitochondrial, Adrenal ferredoxin, Ferredoxin-1, Hepatoredoxin, FDX1, ADX
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	FDX1

Synonyms	ADX
Function	Essential for the synthesis of various steroid hormones (PubMed: 20547883, PubMed:21636783). Participates in the reduction of mitochondrial cytochrome P450 for steroidogenesis (PubMed:20547883, PubMed:21636783). Transfers electrons from adrenodoxin reductase to CYP11A1, a cytochrome P450 that catalyzes cholesterol side-chain cleavage (PubMed:20547883, PubMed:21636783). Does not form a ternary complex with adrenodoxin reductase and CYP11A1 but shuttles between the two enzymes to transfer electrons (By similarity).
Cellular Location	Mitochondrion matrix
Tissue Location	Highest levels in the adrenal gland (at protein level). Also detected in kidney and testis (at protein level)

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat adrenal gland tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Adrenodoxin) Polyclonal Antibody, Unconjugated (AP54496) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.