

CWC22 Rabbit pAb

CWC22 Rabbit pAb

Catalog # AP54513

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9HCG8
Reactivity	Rat, Mouse
Predicted	Pig, Dog, Human, Rabbit, Chicken, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105466
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CWC22
Epitope Specificity	101-200/908
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Spliceosomes are multi-protein complexes that are composed of snRNPs (small nuclear ribonucleoproteins) and a variety of associated protein factors, all of which work in concert to regulate the splicing of pre-mRNA, a critical step in the post-transcriptional regulation of gene expression. CWC22 (CWC22 spliceosome-associated protein), also known as NCM, fSAPb or EIF4GL, is a 908 amino acid nuclear protein and component of the spliceosome C complex. CWC22 is associated with the spliceosome prior to catalytic steps and remains associated throughout the reaction. Belonging to the CWC22 family, CW22 contains one MI domain and a MIF4G domain. The gene encoding CWC22 maps to human chromosome 2q31.3. As the second largest human chromosome, chromosome 2 makes up approximately 8% of the human genome and contains 237 million bases encoding over 1,400 genes.

Additional Information

Gene ID	57703
Other Names	Pre-mRNA-splicing factor CWC22 homolog, Nucampholin homolog, fSAPb, CWC22, KIAA1604, NCM
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	CWC22
Synonyms	KIAA1604, NCM
Function	Required for pre-mRNA splicing as component of the spliceosome (PubMed: 11991638 , PubMed: 12226669 , PubMed: 22961380 , PubMed: 28076346 , PubMed: 28502770 , PubMed: 29301961 , PubMed: 29360106). As a component of the minor spliceosome, involved in the splicing of U12- type introns in pre-mRNAs (Probable). Promotes exon-junction complex (EJC) assembly (PubMed: 22959432 , PubMed: 22961380). Hinders EIF4A3 from non-specifically binding RNA and escorts it to the splicing machinery to promote EJC assembly on mature mRNAs. Through its role in EJC assembly, required for nonsense-mediated mRNA decay.
Cellular Location	Nucleus. Nucleus speckle. Note=Concentrates around speckles, which are sites of pre-mRNA synthesis and processing, where it colocalizes with EJC core proteins.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CWC22) Polyclonal Antibody, Unconjugated (AP54513) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.