

# RSPH4A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54516

## **Product Information**

**Application** WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession Q5TD94

**Reactivity** Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 80733
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human RSPH4A/RSHL3

Epitope Specificity 435-482/716

**Isotype** IgG

**Purity** affinity purified by Protein A

Buffer SUBCELLULAR LOCATION

SIMILARITY
DISEASE

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Cytoplasm; cytoskeleton; cilium axoneme. Radial spoke. Belongs to the flagellar radial spoke RSP4/6 family.

Defects in RSPH4A are the cause of primary ciliary dyskinesia type 11 (CILD11) [MIM:612649]. CILD is an autosomal recessive disorder characterized by axonemal abnormalities of motile cilia. Respiratory infections leading to chronic inflammation and bronchiectasis are recurrent, due to defects in the respiratory cilia; reduced fertility is often observed in male patients due to abnormalities of sperm tails. Half of the patients exhibit situs inversus, due to dysfunction of monocilia at the embryonic node and randomization of left-right body asymmetry. Primary ciliary dyskinesia associated with situs

inversus is referred to as Kartagener syndrome.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** RSHL3 is predicted to be a component of the radial spoke head based on

homology with proteins in the biflagellate alga Chlamydomonas reinhardtii and other ciliates. RSHL3 (radial spoke head-like protein 3), also known as radial spoke head protein 4 homolog A, is a 716 amino acid protein that belongs to the flagellar radial spoke RSP4/6 family. Mutations in the RSHL3 gene cause primary ciliary dyskinesia 1, a disease arising from dysmotility of motile cilia and sperm. Existing as three alternatively spliced isoforms, the RSHL3 gene contains 6 exons, is conserved in chimpanzee, dog, cow, mouse,

rat, chicken, zebrafish, fruit fly and P.falciparum, and maps to human

chromosome 6q22.1.

## **Additional Information**

**Gene ID** 345895

Other Names Radial spoke head protein 4 homolog A, Radial spoke head-like protein 3,

#### RSPH4A, RSHL3

**Target/Specificity** Defects in RSPH4A are the cause of primary ciliary dyskinesia type 11 (CILD11)

[MIM:612649]. CILD is an autosomal recessive disorder characterized by axonemal abnormalities of motile cilia. Respiratory infections leading to chronic inflammation and bronchiectasis are recurrent, due to defects in the respiratory cilia; reduced fertility is often observed in male patients due to abnormalities of sperm tails. Half of the patients exhibit situs inversus, due to dysfunction of monocilia at the embryonic node and randomization of left-right body asymmetry. Primary ciliary dyskinesia associated with situs

inversus is referred to as Kartagener syndrome.

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Protein Information**

Name RSPH4A

Synonyms RSHL3

**Function** Component of the axonemal radial spoke head which plays an important

role in ciliary motility (PubMed: 19200523). Essential for triplet radial spokes


(RS1, RS2 and RS3) head assembly in the motile cilia (By similarity).

**Cellular Location** Cytoplasm, cytoskeleton, cilium axoneme. Cell projection, cilium

**Tissue Location** Expressed in trachea, lungs, and testes (PubMed:23993197). Very strong

expression is detected in nasal brushings (PubMed:19200523).

# **Images**



#### Sample:

Lung (Mouse) Lysate at 40 ug A431(Human) Cell Lysate at 30 ug Testis (Mouse) Lysate at 40 ug

Primary: Anti-RSPH4A (AP54516) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at


1/20000 dilution

Predicted band size: 81 kD Observed band size: 81 kD

### Sample:

Trachea (Mouse) Lysate at 40 ug

Primary: Anti-RSPH4A (AP54516) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution



Predicted band size: 81 kD Observed band size: 81 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.