

SEMA4F Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54518

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O95754
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83511
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SEMA4F
Epitope Specificity	239-288/770
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein
SIMILARITY	Belongs to the semaphorin family.Contains 1 Ig-like C2-type (immunoglobulin-like) domain.Contains 1 PSI domain.Contains 1 Sema domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Semaphorins are a family of cell surface and secreted proteins that are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular "semaphorin" domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA4F (semaphorin-4F), also known as SEMAM, SEMAW or PRO2353, is a 770 amino acid member of the semaphorin family. Localized to the membrane, SEMA4F is a single-pass type I protein that is involved in growth cone collapse of retinal ganglion-cell axons. SEMA4F is highly expressed in postnatal brain and lung and contains one immunoglobulin-like (Ig-like) domain, one PSI domain and one semaphorin domain. Two isoforms exist due to alternative splicing events.

Additional Information

Gene ID	10505
Other Names	Semaphorin-4F, Semaphorin-M, Sema M, Semaphorin-W, Sema W, SEMA4F, SEMAM, SEMAW
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SEMA4F
Synonyms	SEMAM, SEMAW
Function	Probable cell surface receptor that regulates oligodendroglial precursor cell migration (By similarity). Might also regulate differentiation of oligodendroglial precursor cells (By similarity). Has growth cone collapse activity against retinal ganglion-cell axons (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Postsynaptic density {ECO:0000250 UniProtKB:Q9Z143}. Perikaryon {ECO:0000250 UniProtKB:Q9Z123}. Cell projection, dendrite {ECO:0000250 UniProtKB:Q9Z123}. Note=Colocalizes with DLG4 at synapses {ECO:0000250 UniProtKB:Q9Z123}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.