

SEMA6C Rabbit pAb

SEMA6C Rabbit pAb

Catalog # AP54519

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9H3T2
Predicted	Rat, Pig, Dog, Human, Mouse, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	99686
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SEMA6C
Epitope Specificity	256-305/930
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Semaphorins are a family of cell surface and secreted proteins involved in neural development that are conserved from insects to humans. Members of this family are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular "semaphorin" domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. The transmembrane semaphorins are characterized by an additional 80 amino acid transmembrane domain and an 80-110 amino acid cytoplasmic domain. SEMA6C, also known as SEMA Y, is a transmembrane protein expressed in fetal brain and adult skeletal muscle. Three isoforms of this semaphorin exist due to alternative splicing: SEMA6C 1, SEMA6C 2 and SEMA6C 3. The extracellular domain of SEMA6C induces growth cone collapse of dorsal root ganglion and plays a role in generation or stability of entorhino-hippocampal synapses.

Additional Information

Gene ID	10500
Other Names	Semaphorin-6C, Semaphorin-Y, Sema Y, SEMA6C, KIAA1869, SEMAY
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SEMA6C
Synonyms	KIAA1869, SEMAY
Function	Shows growth cone collapsing activity on dorsal root ganglion (DRG) neurons in vitro. May be a stop signal for the DRG neurons in their target areas, and possibly also for other neurons. May also be involved in the maintenance and remodeling of neuronal connections.
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	In adult tissues, expressed only in skeletal muscle.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.