10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# NHLH2 Rabbit pAb

NHLH2 Rabbit pAb
Catalog # AP54532

## Product Information

| | |
|---|---|
| **Application** | WB |
| **Primary Accession** | Q02577 |
| **Reactivity** | Human |
| **Predicted** | Rat, Pig, Dog, Mouse |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 15018 |
| **Physical State** | Liquid |
| **Immunogen** | KLH conjugated synthetic peptide derived from human NHLH2 |
| **Epitope Specificity** | 1-50/135 |
| **Isotype** | IgG |
| **Purity** | affinity purified by Protein A |
| **Buffer** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| **SUBCELLULAR LOCATION** | Nucleus |
| **Important Note** | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |
| **Background Descriptions** | The helix-loop-helix (HLH) structures are known motifs commonly found in membrane-active and DNA-binding proteins. The helix-loop-helix proteins HEN1 and HEN2 are DNA-binding proteins that may be involved in cell-type determination in the early nervous system. Studies of expression in normal tissues have demonstrated expression of NHLH1/NSCL-1 and NHLH2/NSCL-2, the genes encoding HEN1 and HEN2, in the developing central and peripheral nervous system, specifically in developing neurons. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 4808 |
| **Other Names** | Helix-loop-helix protein 2, HEN-2, Class A basic helix-loop-helix protein 34, bHLHa34, Nescient helix loop helix 2, NSCL-2, NHLH2, BHLHA34, HEN2, KIAA0490 |
| **Dilution** | WB=1:500-2000 |
| **Storage** | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |


## Protein Information

| | |
|---|---|
| **Name** | NHLH2 |
| **Synonyms** | BHLHA34, HEN2, KIAA0490 |
| **Function** | Transcription factor which binds the E box motif 5'- CA[TC][AG]TG-3'. Involved in regulating energy expenditure, body mass, voluntary physical activity, mating behavior and reproductive longevity, acting through the hypothalamic-pituitary-gonadal axis. Acts as a transcriptional activator of target genes, including NDN, PCSK1, MC4R (By similarity). Is also a transcriptional activator of KISS1 (PubMed:35066646). May act centrally to regulate function of both white and brown adipose tissue. Together with NHLH1, required to maintain migration and survival of cells in the anterior extramural migration stream (aes), which forms the precerebellar nuclei. Also, in concert with NHLH1, may determine fate of gonadotropin releasing hormone-1 (GnRH-1) neurons. |
| **Cellular Location** | Nucleus {ECO:0000255\|PROSITE-ProRule:PRU00981}. |

## Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## Images




Sample:
PC-3 (human)cell Lysate at 40 ug
Lovo (human)cell Lysate at 40 ug
Primary: Anti- NHLH2 (AP54532) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 15 kD
Observed band size: 18 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.