10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GALP Rabbit pAb

GALP Rabbit pAb
Catalog # AP54533

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UBC7
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	12545
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GALP
Epitope Specificity	25-84/116
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the galanin family of neuropeptides. The encoded protein binds galanin receptors 1, 2 and 3 with the highest affinity for galanin receptor 3 and has been implicated in biological processes involving the central nervous system including hypothalamic regulation of metabolism and reproduction. A peptide encoded by a splice variant of this gene, termed alarin, may have vasoactive properties and serve as a marker for neuroblastic tumors.[provided by RefSeq, Mar 2009].

Additional Information

Gene ID	85569
Other Names	Galanin-like peptide, GALP
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GALP

Function	[Isoform 1]: Hypothalamic neuropeptide which binds to the G-protein-coupled galanin receptors (GALR1, GALR2 and GALR3). Involved in a large number of putative physiological functions in CNS homeostatic processes, including the regulation of gonadotropin-releasing hormone secretion.
Cellular Location	Secreted.
Tissue Location	Isoform 2 is found in ganglia of ganglioneuroma and ganglioneuroblastoma, as well as in differentiated tumor cells of neuroblastoma tissues. Not found in undifferentiated neuroblasts Isoform 2 is found in the skin, in pericytes covering microvascular arterioles and venules on their abluminal surfaces. In larger vessels, isoform 2 is expressed in layers of smooth muscle cells. Isoform 2 is not detected in endothelial cells

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.