

# Gemin 2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54545

#### **Product Information**

**Application** WB, IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Rat, Pig
Host
Clonality
Polyclonal
Calculated MW
31585
Physical State
Liquid

Immunogen KLH conjugated synthetic peptide derived from human Gemin 2

Epitope Specificity 101-200/280

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Nucleus, gem. Cytoplasm. Note=Localized in subnuclear structures next to

coiled bodies, called gems, which are highly enriched in spliceosomal snRNPs.

Also found in the cytoplasm.

**SIMILARITY** Belongs to the gemin-2 family.

**SUBUNIT** Part of the core SMN complex that contains SMN1, GEMIN2/SIP1,

DDX20/GEMIN3, GEMIN4, GEMIN5, GEMIN6, GEMIN7, GEMIN8 and

STRAP/UNRIP. Interacts directly with GEMIN5.

**Important Note**This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** This gene encodes one of the proteins found in the SMN complex, which

consists of several gemin proteins and the protein known as the survival of motor neuron protein. The SMN complex is localized to a subnuclear compartment called gems (gemini of coiled bodies) and is required for assembly of spliceosomal snRNPs and for pre-mRNA splicing. This protein interacts directly with the survival of motor neuron protein and it is required for formation of the SMN complex. A knockout mouse targeting the mouse homolog of this gene exhibited disrupted snRNP assembly and motor neuron

degeneration. [provided by RefSeq, Aug 2011].

### **Additional Information**

Gene ID 8487

Other Names Gem-associated protein 2, Gemin-2, Component of gems 2, Survival of motor

neuron protein-interacting protein 1, SMN-interacting protein 1, GEMIN2,

SIP1

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000

-10000

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

#### **Protein Information**

Name GEMIN2 ( HGNC:10884)

Synonyms SIP1

**Function** The SMN complex catalyzes the assembly of small nuclear


ribonucleoproteins (snRNPs), the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre- mRNAs (PubMed:18984161, PubMed:9323129). Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP (Sm core) (PubMed: 18984161). In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG (5Sm) are trapped in an inactive 6S pICln-Sm complex by the chaperone CLNS1A that controls the assembly of the core snRNP (PubMed:18984161). To assemble core snRNPs, the SMN complex accepts the trapped 5Sm proteins from CLNS1A (PubMed:18984161, PubMed:9323129). Binding of snRNA inside 5Sm ultimately triggers eviction of the SMN complex, thereby allowing binding of SNRPD3 and SNRPB to complete assembly of the core snRNP (PubMed:31799625). Within the SMN complex, GEMIN2 constrains the conformation of 5Sm, thereby promoting 5Sm binding to snRNA containing the snRNP code (a nonameric Sm site and a 3'-adjacent stem-loop), thus preventing progression of assembly until a cognate substrate is bound (PubMed:16314521, PubMed:21816274, PubMed:31799625).

(PubMed:<u>16314521</u>, PubMed:<u>21816274</u>, PubMed:<u>31799625</u>)

**Cellular Location**Nucleus, gem. Cytoplasm. Note=Localized in subnuclear structures next to coiled bodies, called gems, which are highly enriched in spliceosomal snRNPs.

Also found in the cytoplasm

## **Images**



#### Sample:

HepG2(Human) Cell Lysate at 30 ug MCF-7(Human) Cell Lysate at 30 ug

Primary: Anti-Gemin 2 (AP54545) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at


1/20000 dilution

Predicted band size: 32 kD Observed band size: 32 kD


Paraformaldehyde-fixed, paraffin embedded (rat spinal cord); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Gemin 2) Polyclonal Antibody, Unconjugated (AP54545) at 1:200 overnight at 4°C,



followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat thymus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Gemin 2) Polyclonal Antibody, Unconjugated (AP54545) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Gemin 2) Polyclonal Antibody, Unconjugated (AP54545) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.