

Nicalin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54561

Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession

Reactivity

Rat

Host

Clonality

Polyclonal

Calculated MW

Ogeonal

Additional Information

Gene ID 56926

Other Names Nicalin, Nicastrin-like protein, NCLN

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name NCLN {ECO:0000303 | PubMed:36261522, ECO:0000312 | HGNC:HGNC:26923}

FunctionComponent of the multi-pass translocon (MPT) complex that mediates insertion of multi-pass membrane proteins into the lipid bilayer of

membranes (PubMed:32820719, PubMed:36261522). The MPT complex takes over after the SEC61 complex: following membrane insertion of the first few transmembrane segments of proteins by the SEC61 complex, the MPT

complex occludes the lateral gate of the SEC61 complex to promote insertion of subsequent transmembrane regions (PubMed:36261522). May antagonize Nodal signaling and subsequent organization of axial structures during mesodermal patterning, via its interaction with NOMO (By similarity).

Cellular Location Endoplasmic reticulum membrane; Single-pass membrane protein

Tissue Location Highly expressed in pancreas and skeletal muscle and, at lower levels, in

heart.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.