

UBXD2 Rabbit pAb

UBXD2 Rabbit pAb
Catalog # AP54592

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q92575
Reactivity	Rat, Human, Mouse
Predicted	Pig, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56778
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human UBXD2
Epitope Specificity	201-300/508
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum membrane. Both the N- and the C-terminus face the cytosol; also found in the nucleus envelope contiguous to the ER.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Erasin is an endoplasmic reticulum (ER) and nuclear envelope membrane protein. Expressed in a variety of tissues, such as brain, placenta, heart, liver, prostate, kidney, pancreas, lung and skeletal muscle, erasin contains one UBX domain and participates in the clearing of ERAD (endoplasmic reticulum-associated protein degradation) substrates. The UBX domain of erasin is responsible for mediating its direct interaction with VCP (valosin-containing protein), an AAA-ATPase molecular chaperone. In response to ER stress, erasin expression is induced. The knockdown of erasin expression leads to the inhibition of ERAD, suggesting an important function of erasin in the ERAD pathway. In addition, erasin may be involved in Alzheimer's disease, as it is known to accumulate in neurofibrillary degenerating neurons in patients with Alzheimer's disease.

Additional Information

Gene ID	23190
Other Names	UBX domain-containing protein 4, Erasin, UBX domain-containing protein 2, UBXN4, KIAA0242, UBXD2, UBXDC1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	UBXN4
Synonyms	KIAA0242, UBXD2, UBXDC1
Function	Involved in endoplasmic reticulum-associated protein degradation (ERAD). Acts as a platform to recruit both UBQLN1 and VCP to the ER during ERAD (PubMed: 19822669).
Cellular Location	Endoplasmic reticulum membrane; Peripheral membrane protein. Nucleus envelope. Note=Both the N- and the C-terminus face the cytosol. Also found in the nucleus envelope contiguous to the ER
Tissue Location	Expressed in many tissues, including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas Accumulates in Alzheimer disease-afflicted brains (at protein level)

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:

Heart (Mouse) Lysate at 40 ug


Cerebrum (Rat) Lysate at 40 ug

Primary: Anti-UBXD2 (AP54592) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 57 kD

Observed band size: 57 kD



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UBXD2) Polyclonal Antibody, Unconjugated (AP54592) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.