

# Capicua Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54608

## Product Information

|                                |   |
|--------------------------------|---|
| <b>Application</b>             | IHC-P, IHC-F, IF, ICC, E  |
| <b>Primary Accession</b>       | <a href="#">Q96RK0</a>  |
| <b>Reactivity</b>              | Rat, Pig, Dog, Bovine   |
| <b>Host</b>                    | Rabbit  |
| <b>Clonality</b>               | Polyclonal  |
| <b>Calculated MW</b>           | 258033  |
| <b>Physical State</b>          | Liquid  |
| <b>Immunogen</b>               | KLH conjugated synthetic peptide derived from human Capicua   |
| <b>Epitope Specificity</b>     | 1475-1560/1608  |
| <b>Isotype</b>                 | IgG   |
| <b>Purity</b>                  | affinity purified by Protein A  |
| <b>Buffer</b>                  | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.   |
| <b>SUBCELLULAR LOCATION</b>    | Nucleus.  |
| <b>SIMILARITY</b>              | Contains 1 HMG box DNA-binding domain.  |
| <b>SUBUNIT</b>                 | Interacts with ATXN1 and ATXN1L   |
| <b>Important Note</b>          | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.   |
| <b>Background Descriptions</b> | Capicua is the mammalian ortholog of the drosophila Cic gene and is part of the HMG-box protein superfamily. Expressed primarily in the fetal brain, Capicua functions as a transcriptional repressor and is involved in the development of the nervous system through interaction with the ATXN1 protein. When ATXN1 assembles into stable complexes, it directly binds Capicua, thereby mediating both the activity and expression of Capicua. When Capicua is active, it is able to interact with other developmental proteins to restrict the growth of granule cells and regulate normal neuronal development. Disruptions in the the association of Capicua with proteins such as ATXN1 are thought to cause medulloblastoma, the most common form of peridiatric brain tumor arising from irregular growth of granule cells. |

## Additional Information

|                           |   |
|---------------------------|---|
| <b>Gene ID</b>            | 23152   |
| <b>Other Names</b>        | Protein capicua homolog, CIC, KIAA0306  |
| <b>Target/Specificity</b> | Expressed in fetal brain.   |
| <b>Dilution</b>           | IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000 |
| <b>Format</b>             | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce          |

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


## Protein Information

---

|                          |   |
|--------------------------|---|
| <b>Name</b>              | CIC   |
| <b>Synonyms</b>          | KIAA0306  |
| <b>Function</b>          | Transcriptional repressor which plays a role in development of the central nervous system (CNS). In concert with ATXN1 and ATXN1L, involved in brain development. |
| <b>Cellular Location</b> | Nucleus {ECO:0000255 PROSITE-ProRule:PRU00267}.   |
| <b>Tissue Location</b>   | Expressed in fetal brain.   |

## Images

---



Tissue/cell: brain of mouse embryo; 4% Paraformaldehyde-fixed and paraffin-embedded;  
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;  
Incubation: Anti-Capicua Polyclonal Antibody, Unconjugated(AP54608) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.