

CSDC2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54637

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9Y534
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16786

Additional Information

Gene ID	27254
Other Names	Cold shock domain-containing protein C2, RNA-binding protein PIPPIN, CSDC2, PIPPIN
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CSDC2
Synonyms	PIPPIN
Function	RNA-binding factor which binds specifically to the very 3'-UTR ends of both histone H1 and H3.3 mRNAs, encompassing the polyadenylation signal. Might play a central role in the negative regulation of histone variant synthesis in the developing brain (By similarity).
Cellular Location	Nucleus. Cytoplasm. Note=PIPPin-RNA complexes are located to the nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.