

TREM2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5469A

Product Information

Application	WB, E
Primary Accession	Q9NZC2
Other Accession	NP_061838.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB26747
Calculated MW	25447
Antigen Region	22-50

Additional Information

Gene ID	54209
Other Names	Triggering receptor expressed on myeloid cells 2, TREM-2, Triggering receptor expressed on monocytes 2, TREM2
Target/Specificity	This TREM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between a range from 22-50 amino acids of human TREM2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TREM2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TREM2 (HGNC:17761)
Function	Forms a receptor signaling complex with TYROBP which mediates signaling and cell activation following ligand binding (PubMed: 10799849). Acts as a receptor for amyloid-beta protein 42, a cleavage product of the amyloid-beta

precursor protein APP, and mediates its uptake and degradation by microglia (PubMed:[27477018](#), PubMed:[29518356](#)). Binding to amyloid-beta 42 mediates microglial activation, proliferation, migration, apoptosis and expression of pro- inflammatory cytokines, such as IL6R and CCL3, and the anti-inflammatory cytokine ARG1 (By similarity). Acts as a receptor for lipoprotein particles such as LDL, VLDL, and HDL and for apolipoproteins such as APOA1, APOA2, APOB, APOE, APOE2, APOE3, APOE4, and CLU and enhances their uptake in microglia (PubMed:[27477018](#)). Binds phospholipids (preferably anionic lipids) such as phosphatidylserine, phosphatidylethanolamine, phosphatidylglycerol and sphingomyelin (PubMed:[29794134](#)). Regulates microglial proliferation by acting as an upstream regulator of the Wnt/beta-catenin signaling cascade (By similarity). Required for microglial phagocytosis of apoptotic neurons (PubMed:[24990881](#)). Also required for microglial activation and phagocytosis of myelin debris after neuronal injury and of neuronal synapses during synapse elimination in the developing brain (By similarity). Regulates microglial chemotaxis and process outgrowth, and also the microglial response to oxidative stress and lipopolysaccharide (By similarity). It suppresses PI3K and NF-kappa-B signaling in response to lipopolysaccharide; thus promoting phagocytosis, suppressing pro-inflammatory cytokine and nitric oxide production, inhibiting apoptosis and increasing expression of IL10 and TGFB (By similarity). During oxidative stress, it promotes anti-apoptotic NF- kappa-B signaling and ERK signaling (By similarity). Plays a role in microglial MTOR activation and metabolism (By similarity). Regulates age-related changes in microglial numbers (PubMed:[29752066](#)). Triggers activation of the immune responses in macrophages and dendritic cells (PubMed:[10799849](#)). Mediates cytokine-induced formation of multinucleated giant cells which are formed by the fusion of macrophages (By similarity). In dendritic cells, receptor of SEMA6D with PLEXNA1 as coreceptor and mediates up-regulation of chemokine receptor CCR7 and dendritic cell maturation and survival (PubMed:[11602640](#)). Involved in the positive regulation of osteoclast differentiation (PubMed:[12925681](#)).

Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Secreted.
Tissue Location	Expressed in the brain, specifically in microglia and in the fusiform gyrus (at protein level) (PubMed: 27477018 , PubMed: 28802038 , PubMed: 28855300 , PubMed: 29752066). Expressed on macrophages and dendritic cells but not on granulocytes or monocytes (PubMed: 10799849 , PubMed: 28855301). In the CNS strongest expression seen in the basal ganglia, corpus callosum, medulla oblongata and spinal cord (PubMed: 12080485).


Background

TREM2 is a membrane protein that forms a receptor signaling complex with TYROBP. The encoded protein may be involved in chronic inflammation by triggering the production of constitutive inflammatory cytokines. Defects in this gene are a cause of polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOS).

References

Stefano, L., et al. J. Neurochem. 110(1):284-294(2009)
 Ford, J.W., et al. Curr. Opin. Immunol. 21(1):38-46(2009)

Images



All lanes: Anti-TREM2Antibody(N-term) at 1:1000 dilution
Lane 1: SH-SY5Y whole cell lysate Lane 2: Mouse lung cell lysate
Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 27 KDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.