

GPRC5B Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54708

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q9NZH0

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 44795
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human G protein coupled

receptor family C group 1 member B

Epitope Specificity 1-100/403

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane. Cytoplasmic vesicle membrane. Localized in the plasma

membrane and perinuclear vesicles.

SIMILARITY Belongs to the G-protein coupled receptor 3 family.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions GPRC5B (G protein-coupled receptor family C group 5 member B, retinoic

acid-induced gene 2 protein) is a 403 amino acid protein encoded by the human GPRC5B gene. GPRC5B is an orphan receptor member of the G protein-coupled receptor 3 family. G protein-coupled receptors (GPCRs or GPRs) contain 7 transmembrane domains and transduce extracellular signals through heterotrimeric G proteins. Key roles for G protein-coupled receptors include control of protein maturation and cell surface delivery, and providing the correct framework for interactions with both heterotrimeric G proteins and arrestins to allow signal generation and termination. This retinoic acid-inducible G protein-coupled receptor provides evidence for a possible interaction between retinoid and G protein signaling pathways. GPRC5B is highly expressed in kidney, pancreas and testis, and has moderate expression

in brain, heart, prostate, small intestine and spleen.

Additional Information

Gene ID 51704

Other Names G-protein coupled receptor family C group 5 member B, A-69G12.1, Retinoic

acid-induced gene 2 protein, RAIG-2, GPRC5B, RAIG2

Target/Specificity Expression is high in kidney, pancreas, and testis, medium in brain, heart,

prostate, small intestine, and spleen, low in liver, placenta, skeletal muscle,

colon, ovary, and thymus, and not detectable in lung and peripheral leukocyte. According to PubMed:10945465, highly expressed in most brain areas examined, with the highest levels observed in corpus callosum, caudate nucleus, putamen, substantia nigra, thalamus, hippocampus, and spinal chord as well as in dorsal root ganglia (DRG). In the periphery, expression levels are relatively low, compared to the CNS, with the strongest expression detected in pancreas, testis, uterus, and stomach.

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,Flow-Cyt=1

©g/Test,ELISA=1:5000-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name GPRC5B

Synonyms RAIG2

Function G-protein coupled receptor involved in the regulation of cell volume.

Cellular Location Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle

membrane; Multi-pass membrane protein. Note=Localized in the plasma


membrane and perinuclear vesicles

Tissue Location Expression is high in kidney, pancreas, and testis, medium in brain, heart,

prostate, small intestine, and spleen, low in liver, placenta, skeletal muscle, colon, ovary, and thymus, and not detectable in lung and peripheral leukocyte. According to PubMed:10945465, highly expressed in most brain areas examined, with the highest levels observed in corpus callosum, caudate nucleus, putamen, substantia nigra, thalamus, hippocampus, and spinal cord as well as in dorsal root ganglia (DRG). Expressed in glia limitans, ependymal cells, astrocyte cell bodies, the perivascular region in astrocyte endfeet, but not in neurons (PubMed:37143309). In the periphery, expression levels are relatively low, compared to the CNS, with the strongest expression detected in

pancreas, testis, uterus, and stomach.

Images



Positive control: (mo)Splenocytes(2%

Paraformaldehyde-fixed)

Isotype Control Antibody: Rabbit IgG; Dilution: 1 µg in

100 µl 1 X PBS containing 0.5% BSA

Secondary Antibody: Goat anti-rabbit IgG-FITC; Dilution:

1:200 in 1 X PBS containing 0.5% BSA

Primary Antibody: rabbit Anti-GPRC5B (AP54708); Dilution: 1 µg in 100 µl 1X PBS containing 0.5% BSA

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.