10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

5HT7 Receptor Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54714

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	P34969
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53555

Additional Information

Gene ID	3363
Other Names	5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT7, 5-HT-X, Serotonin receptor 7, HTR7
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HTR7 {ECO:0000303|Ref.3, ECO:0000312|HGNC:HGNC:5302}
Function	G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed:35714614, PubMed:8226867). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed:35714614, PubMed:8226867). HTR7 is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed:35714614).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	[Isoform A]: Predominant isoform in spleen, caudate and hippocampus. [Isoform D]: Minor isoform in terms of expression.

Images



Sample:
Hippocampus (Mouse) Lysate at 40 ug
Hippocampus(Rat) Lysate at 40 ug
Primary: Anti- 5HT7 Receptor (AP54714) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 54 kD
Observed band size: 54 kD