

FAM62C Rabbit pAb

FAM62C Rabbit pAb

Catalog # AP54757

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	A0FGR9
Reactivity	Rat
Predicted	Pig, Dog, Human, Mouse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	100035
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FAM62C
Epitope Specificity	101-200/886
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Single-pass membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 3 is made up of about 214 million bases encoding over 1,100 genes. Notably, there is a chemokine receptor gene cluster and a variety of human cancer related loci on chromosome 3. Particular regions of the chromosome 3 short arm are deleted in many types of cancer cells. Key tumor suppressing genes on chromosome 3 encode apoptosis mediator RASSF1, cell migration regulator HYAL1 and angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth Disease are a few of the numerous genetic diseases associated with chromosome 3. The FAM62C gene product has been provisionally designated FAM62C pending further characterization.

Additional Information

Gene ID	83850
Other Names	Extended synaptotagmin-3, E-Syt3, Chr3Syt, ESYT3 (HGNC:24295), FAM62C
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ESYT3 (HGNC:24295)
Synonyms	FAM62C
Function	Binds glycerophospholipids in a barrel-like domain and may play a role in cellular lipid transport (By similarity). Tethers the endoplasmic reticulum to the cell membrane and promotes the formation of appositions between the endoplasmic reticulum and the cell membrane.
Cellular Location	Cell membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Note=Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS) (PubMed:29469807, PubMed:30220461). Recruited to the cell membrane via the third C2 domain
Tissue Location	Widely expressed with high level in cerebellum and skin.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM62C) Polyclonal Antibody, Unconjugated (AP54757) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.